

# the ADHD REPORT

Russell A. Barkley & Associates

• Volume 26

• Number 5

• ISSN 1065-8025

• August 2018

## Physical Exercise Interventions for Emerging Adults with Attention-Deficit/Hyperactivity Disorder (ADHD)

Patrick A. LaCount, M.S., and Cynthia M. Hartung, Ph.D.

Attention-deficit/hyperactivity disorder (ADHD) is characterized by deficits in executive functioning (e.g., working memory, response inhibition, and organization; Willcutt, Doyle, Nigg, Faraone, & Pennington, 2005). Inattention and hyperactivity/impulsivity result in impairments in academic, occupational, and social functioning (American Psychiatric Association [APA], 2013). The primary treatments for ADHD include psychosocial, pharmacological, and combined treatments. Despite the well-documented effectiveness of these interventions, there are limitations to their use. Although both pharmacotherapy and psychosocial treatments are efficacious during active treatment (Knight, Rooney, & Chronis-Tuscano, 2008; Spencer et al., 1996), few individuals receive long-term treatment, resulting in limited, if any, sustained effects (Molina et al., 2009). One explanation for the transient benefits of current treatments is that they target symptoms and functional impairments rather than the neural mechanisms that underlie the disorder (Berwid & Halperin, 2012). In response to this, re-

searchers have begun to consider other forms of interventions to directly impact these underlying deficits.

Physical exercise has received increasing attention with the current obesity epidemic (e.g., Davis et al., 2011). The benefits of physical exercise include improving medical problems (e.g., obesity, diabetes) and psychological difficulties (e.g., anxiety, depression; Hillman, Erickson, & Kramer, 2008). A growing body of research suggests physical exercise has powerful ef-

fects on neurocognitive, psychological, and academic functioning (Hillman et al., 2008; Trost, Owen, Bauman, Sallis, & Brown, 2002). Emerging from this burgeoning evidence, mental health researchers have begun to explore exercise as an avenue by which mental health treatment outcomes can be enhanced.

Among individuals with ADHD, exercise may not only improve cognitive performance (Berwid & Halperin, 2012; Gapin, Labban, & Etnier, 2011), but

### Contents

**Physical Exercise Interventions for Emerging Adults with Attention-Deficit/Hyperactivity Disorder (ADHD), 1 •**

**Research Findings, 12 •**

NOTICE TO NON-PROFESSIONALS The information contained in this newsletter is not intended as a substitute for consultation with health care professionals.

Russell A. Barkley, Ph.D.  
Virginia Treatment Center for Children  
Virginia Commonwealth University  
School of Medicine  
Send correspondence to  
drbarkley@russellbarkley.org

Dr. Russell Barkley discloses that his work includes serving as a paid Speaker/Consultant for pharmaceutical companies in his area of expertise.

#### ADVISORY BOARD

Kevin Antshel, Ph.D., Syracuse University • José J. Bauermeister, Ph.D., University of Puerto Rico, San Juan • Stephen P. Becker, Ph.D., Cincinnati Children's Hospital Medical Center • Will Canu, Ph.D., Appalachian State University • Anil Chacko, Ph.D., New York University • Andrea M. Chronis, Ph.D., University of Maryland, College Park • George J. DuPaul, Ph.D., Lehigh University, PA • Gregory Fabiano, Ph.D., University of Buffalo • Jeff Epstein, Ph.D., Cincinnati Children's Hospital Medical Center • Sam Goldstein, Ph.D., University of Utah, Salt Lake City • Cynthia M. Hartung, Ph.D., University of Wyoming • Stephen Hinshaw, Ph.D., UC Berkeley • Betsy Hoza, Ph.D., University of Vermont • Charlotte Johnston, Ph.D., University of British Columbia, Vancouver • Laura E. Knouse, Ph.D., University of Richmond, VA • Scott Kollins, Ph.D., Duke University Medical Center, Durham, NC • Sandra Kooij, M.D., Expertise Center Adult ADHD, The Netherlands • Joshua Langberg, Ph.D., Virginia Commonwealth University • Florence Levy, M.D., The Prince of Wales Children's Hospital, Australia • Larry Lewandowski, Ph.D., Syracuse University, NY • Sandra Loo, Ph.D., Neuropsychiatric Institute, UCLA • Keith McBurnett, Ph.D., University of California, San Francisco • Richard Milich, Ph.D., University of Kentucky, Lexington • Brooke Molina, Ph.D., University of Pittsburgh • Kevin Murphy, Ph.D., Adult ADHD Clinic of Central Massachusetts • Joel Nigg, Ph.D., Oregon Health Sciences University, Portland, OR • Linda Pfiffner, Ph.D., University of Chicago • J. Russell Ramsay, Ph.D., University of Pennsylvania Perelman School of Medicine • Mark Rapport, Ph.D., University of Central Florida • Luis Rohde, M.D., Federal University of Rio Grande do Sul • Julie Schweitzer, Ph.D., UC-Davis MIND Institute, Sacramento, CA • Mary V. Solanto, Ph.D., Hofstra Northwell School of Medicine • Dan Waschbusch, Ph.D., Penn State University Medical Center • Jeanette Wasserstein, Ph.D., The Mount Sinai School of Medicine • Lisa Weyandt, Ph.D., University of Rhode Island • Alan Zametkin, M.D., Private Practice

THE ADHD REPORT (ISSN 1065-8025) is published bimonthly by The Guilford Press, 370 Seventh Avenue, Suite 1200, New York, NY 10001-1020. Guilford's GST registration number: 137401014.

Subscription price: (eight issues) Individuals \$105.00, Institutions, \$470.00. Add \$15.00 for Canada and Foreign (includes airmail postage). Orders by MasterCard, VISA, or American Express can be placed by Phone at 800-365-7006, Fax 212-966-6708, or E-mail news@guilford.com; in New York, 212-431-9800. Payment must be made in U.S. dollars through a U.S. bank. All prices quoted in U.S. dollars. Pro forma invoices issued upon request.

Visit our website at [www.guilford.com](http://www.guilford.com).

CHANGE OF ADDRESS: Please inform publisher at least six weeks prior to move. Enclose mailing label with change of address. Claims for lost issues cannot be honored four months after mailing date. Duplicate copies cannot be sent to replace issues not delivered because of failure to notify publisher of change of address. Postmaster: Change of address to The ADHD Report, Guilford Press, 370 Seventh Avenue, Suite 1200, New York, NY 10001-1020.

Photocopying of this newsletter is not permitted.

Inquire for bulk rates.

Copyright © 2018 by The Guilford Press.

Printed in the United States of America.

ameliorate comorbid anxiety and depression (Salmon, 2001), and serve as a protective factor for adverse health outcomes (Nigg, 2013)—all of which commonly co-occur with ADHD (Barkley, Murphy, & Fischer, 2008). Indeed, initial results with children and adolescents have been promising. Findings from these studies provided preliminary evidence that exercise may improve motor, cognitive, social, and behavioral functioning for school-aged children with ADHD (Neudecker, Mewes, Reimers, & Woll, 2015). While investigations on interventions involving exercise for school-aged children with ADHD advance, there are no known studies examining the use of such interventions for ADHD among *emerging adults* (ages 18–25; Arnett, 2000).

Although tempting, before generalizing school-aged treatment research to emerging adults, there are several important developmental and environmental differences that must be considered. For example, emerging adults experience higher demands for self-regulation, have greater autonomy, diminished external support (e.g., parents, schools), and their neurological structures are near their peak maturity (Giedd, 2004). Thus, tailoring school-aged interventions to the developmental and environmental context of emerging adulthood is warranted to evaluate possible treatment effects.

We summarized research across disciplines to elucidate the relation between exercise and ADHD-related impairment in the developmental context of emerging adulthood. We also incorporated research on possible mechanisms of change specific to ADHD and exercise interventions for depression—the mental health area with which the literature is most robust—to understand the strengths and weaknesses of this therapeutic approach outside of childhood. We conclude with clinical implications and recommendations, as well as avenues for future research.

#### PHYSICAL FITNESS, ACTIVITY, AND EXERCISE

A robust literature suggests that physical fitness, physical activity, and physical exercise have powerful effects on the

structure, function, and development of the brain over both the short term and the long term (Best, 2010; Hillman et al., 2008; Portugal et al., 2013; Tomporowski, Lambourne, & Okumura, 2011). However, these terms are often confused with one another and are often used interchangeably even though they describe different concepts. The following section provides definitions and methods for quantifying physical fitness, physical activity, and physical exercise as outlined by the Centers for Disease Control and Prevention (CDC; see Casperson, Powell, & Christenson, 1985).

#### Physical Fitness

Whereas physical activity and physical exercise are related to the movements that people perform, *physical fitness* corresponds to attributes people have or achieve. Being physically fit has been broadly described as the ability to complete daily tasks with vigor and alertness, with ample energy to enjoy leisure pursuits and meet unforeseen emergencies. Many quantifiable components contribute to physical fitness and are categorized as being skill- or health-related fitness.

Skill-related components of physical fitness and can be categorized into six components: (1) agility, (2) balance, (3) coordination, (4) speed, (5) power, and (6) reaction time. There are many methods for measuring each of the six components of skill-related physical fitness, including: weaving through cones, balancing on one leg, throwing and catching a ball, sprints, jumping up on a box, and hitting a thrown baseball with a bat, respectively.

The five health-related components of physical fitness include: (1) cardiovascular endurance, (2) muscular endurance, (3) muscular strength, (4) body composition, and (5) flexibility. Similar to skill-related physical fitness, there are multiple methods for measuring each of the components of health-related physical fitness, including: a 12-minute run, sit-ups, handgrip dynamometer, body mass index, and sit-and-reach test, respectively.

## Physical Activity

*Physical activity* is any bodily movement contrived by skeletal muscles that results in energy expenditure above the basal metabolic rate. This includes, but is not limited to, physical activity occurring at work (e.g., sitting, standing, lifting), leisure (e.g., household tasks, sports, metabolic conditioning exercises), and while sleeping.<sup>1</sup> Energy expenditure is often measured in kilocalories (kcal) or kilojoules (kJ) and can be expressed as a total amount of energy used in a given physical activity and rate (kcal per unit of time). The net amount of energy used with a physical activity is associated with the amount of muscle mass used to produce bodily movements and the intensity, duration, and frequency of muscular contractions.

## Physical Exercise

Although the terms *physical activity* and *physical exercise* have many common elements, these terms are not synonymous. *Physical exercise* is any physical activity that is planned, structured, and repetitive with the intention to develop or maintain physical fitness (e.g., strength, flexibility, aerobic endurance). Therefore, physical exercise is a subtype of physical activity and may compose all or part of different physical activities, except for sleep. For example, the physical activity of playing sports is often purposefully performed to improve or maintain physical fitness and is often planned, structured, and repetitive. Conversely, household and work tasks are often completed in a labor-saving manner. That is, the goals of these physical activities are often completed with the goal of energy conservation, with little to no regard to physical fitness and would therefore be classified as physical activity. However, a person may plan or structure the performance of household or work tasks in a labor-inducing manner to “burn” more calories or develop muscular strength. When tasks are performed in this manner, they are considered physical exercise.

The terms *physical activity* and *physical exercise* are often conflated in the ADHD literature, which obfuscates interpretations of empirical research. This is compounded by the broad range of physical activities/exercises in school-aged ADHD studies, which include aquatic exercise (Chang, Hung, Huang, Hatfield, & Hung, 2014), plyometric exercise (tuck jumps, one-leg hops; McKune, Pautz, & Lomjbard, 2004), 20 minutes of walking (Taylor & Kuo, 2009), recumbent cycling (Piepmeier et al., 2015), yoga (Jensen & Kenney, 2004), and sports (basketball, soccer, tennis; Pan et al., 2015; Verret, Guay, Berthiaume, Gardiner, & Béliveau, 2012). This diversity of methods results in variable intensity, duration, and frequency of interventions across studies, precluding uniform comparison of the effects of physical activity interventions (e.g., walking rather than being driven to school) and physical exercise interventions (e.g., participating in vigorous aerobic activity) for children and adolescents with ADHD. However, because most investigations on this topic use interventions methods that satisfy all or most of the definitional criteria for physical exercise (PE), this term will be used to refer to both physical exercise and physical activity for the remainder of the article.

## NEUROPSYCHOLOGY OF ADHD

Over the last 10 years, several meta-analyses have been performed on different aspects of cognition and neurobiology of ADHD (e.g., Hervey, Epstein, & Curry, 2004; Schoechlin & Engel, 2005; Willcutt et al., 2005; Willcutt et al., 2012; Woods, Lovejoy, & Ball, 2002). The following section is a summary of the cognitive and neurobiological mechanisms implicated in the manifestation of ADHD (for a detailed review, see Willcutt, 2015).

## Cognitive Factors Related to ADHD

Prevailing cognitive models suggest that ADHD arises from general weaknesses in executive functioning (EF)—a set of “top-down” cognitive processes

used to regulate behavior toward adaptive, long-term goals (Barkley, 1997). Specific EF processes implicated in ADHD include the ability to inhibit impulses (inhibition), short-term storage and manipulation of information (working memory), and preferring smaller, immediate rewards over larger, delayed rewards (delay aversion; Willcutt et al., 2012). Across the lifespan, meta-analyses report medium to large differences among groups with and without ADHD across these EF tasks ( $d = .50-.70$ ; Willcutt, 2015). However, the patterns of cognitive impairments observed in ADHD are inconsistent across individuals.

Even though individuals with ADHD are 3 to 4 times more likely to exhibit significant deficits in EF than those without ADHD, less than half of individuals with ADHD exhibit significant deficits on any specific EF tasks (Nigg et al., 2005). Similarly, performance in EF tasks tends to correlate significantly with ADHD symptoms but are small in magnitude ( $r = .15-.35$ ; Nigg, Hinshaw, Carte, & Treuting, 1998; Willcutt et al., 2001). Thus, the cognitive processes underlying ADHD are complex and multifactorial, with no single EF deficit being sufficient or necessary to characterize all cases of ADHD (e.g., Pennington, 2006; Willcutt et al., 2010).

## Neurobiological Factors Related to ADHD

Consistent with the cognitive literature, studies indicate that there is no single identifiable neurobiological etiology for ADHD. However, models of ADHD posit that neuroanatomical and neurochemical abnormalities in the prefrontal cortex (PFC) underlie much of the cognitive and behavioral manifestations of ADHD (Berridge & Devilbiss, 2011; Durston & Konrad, 2007). The PFC is a region of the brain that is frequently cited as being central to the pathophysiology of ADHD—particularly the frontostriatal and mesolimbic networks (Castellanos & Tannock, 2002; Durston, 2003).

1. Although the caloric contribution of physical activity during sleep accounts for a small portion of energy expenditure due to physical activity, the energy used is above basal metabolic rate and is therefore classified as a category of physical activity (Caspersen et al., 1985).

Working memory and inhibition are largely supported by the frontostriatal network, which has links to the cerebellum through the thalamus and is comprised of the lateral prefrontal cortex, dorsal anterior cingulate, and dorsal striatum (Durstun, 2003). Structural neuroimaging research has found that adults with ADHD show cortical thinning in the dorsolateral prefrontal and anterior cingulate cortices (Makris et al., 2007). Further, a meta-analysis of neuroimaging studies found a consistent pattern of hypoactivity in the frontostriatal network among children and adolescents with ADHD during tasks requiring working memory and inhibition (Dickstein, Bannon, Xavier Castellanos, & Milham, 2006).

Delay aversion is supported by the mesolimbic network, which connects the ventral striatum (nucleus accumbens), areas of the hippocampus and amygdala, ventral tegmental area, and ventromedial prefrontal regions, which includes the orbitofrontal gyri and anterior cingulate (Sonuga-Barke, 2002). Across studies examining the impact of reinforcement contingencies on ADHD, those with ADHD tend to show atypical physiological responses to rewards, as well as risky decision-making and greater delay aversion (Luman, Oosterlaan, & Sergeant, 2005). Functional neuroimaging studies have reported that individuals with ADHD show lower activation in the ventral striatum when anticipating reward, but higher activation when they received the reward (Scheres, Milham, Knutson, & Castellanos, 2007; Ströhle et al., 2008). Individuals with ADHD also show lower top-down inhibitory control in the orbitofrontal cortex and ventral striatum (Ströhle et al., 2008). Further, structural neuroimaging research has found that adults with ADHD show cortical thinning in the anterior cingulate cortex (Makris et al., 2007).

Neurotransmitter dysfunction has also been implicated in the neurobiological pathology of ADHD (Arnsten, 2009). Dysregulation of dopamine and norepinephrine in the PFC has been suggested to underlie the EF deficits observed in ADHD (Pliszka, 2005). The dopaminergic system, which is associated with motor control, motivation, re-

ward, and affect, is believed to interact with the PFC as a regulation network that is associated with cognitive and behavioral control (Wigal, Emmerson, Gehricke, & Galassetti, 2012). Experimental studies have shown that working memory tasks increase dopamine levels in the PFC, and obstructing dopamine receptors in the PFC creates working memory impairments (Durstewitz, Kelc, & Güntürkün, 1999). Given that individuals with ADHD often display impairments in working memory, these studies suggest that dysregulation of dopamine may correspond to the cognitive deficits observed in ADHD (Wigal et al., 2012). Specifically, it is hypothesized that attentional deficit may be caused by a hypodopaminergic state in the PFC (Solanto, 2002). Similarly, the noradrenergic system, which is involved in attention, working memory, and inhibition, is thought to be involved in the pathophysiology of ADHD due to low norepinephrine levels in the PFC being associated with deficits in working memory and EF (Arnsten, 2000; Pliszka, 2005). These studies of neurochemical dysfunction in ADHD are further supported by research on ADHD medications, which enhance levels of dopamine and norepinephrine in the PFC and ameliorate ADHD symptomatology (Pliszka, 2005).

### COGNITIVE AND NEUROBIOLOGICAL EFFECTS OF PHYSICAL EXERCISE

Human and non-human animal studies have shown that PE results in short- and long-term neurobiological effects and can improve several aspects of cognitive functioning (for reviews, see Hillman et al., 2008; Loprinzi, Herod, Cardinal, & Noakes, 2013). PE has been hypothesized to be an effective therapeutic tool to target the neurobiological mechanisms associated with ADHD (Berwid & Halperin, 2012; Gapin et al., 2011; Wigal et al., 2012). The following section provides an overview of the cognitive and neurobiological effects of PE with a focus on components that underlie ADHD to identify theoretical mechanisms of change for PE interventions for emerging adults with ADHD.

### COGNITIVE EFFECTS OF PHYSICAL EXERCISE

Several meta-analytic reviews have found positive associations between PE and cognitive functioning across the lifespan (Angevaren et al., 2008; Etner, Nowell, Landers, & Sibley, 2006; Kramer & Erickson, 2007; Sibley & Etner, 2003). However, the relation and improvements related to PE are not homogenous across domains of cognitive functioning. Across studies, PE consistently has been shown to disproportionately improve EF in children and adults (Hillman et al., 2008; Voss, Carr, Clark, & Weng, 2014). However, there is a relative lack of research on PE-induced cognitive effects in emerging adults. This is likely due to emerging adults often serving as comparisons for older adults and being near peak neurodevelopmental maturity (Giedd, 2004; Hillman et al., 2008). Existing literature indicates, however, that PE can yield cognitive benefits in emerging adulthood. A large cohort study demonstrated a positive association between PE and cognitive functioning in emerging adults (Åberg et al., 2009). Although information regarding influences on EF was not available for this study, PE was most strongly associated with logical reasoning, which contains components of EF. Additionally, Sibley and Beilock (2007) found that improvements in working memory were influenced by PE, but only for emerging adults who were impaired in this cognitive domain ( $d = 1.22$ ). This suggests that PE may be uniquely beneficial for emerging adults with impaired cognitive functioning, such as ADHD.

### Neurobiological Effects of Physical Exercise

Engaging in PE has a variety of effects on the brain's structure and functioning, including neurotransmitter and brain-derived neurotrophic factor (BDNF) levels, neurogenesis, synaptic plasticity, neuroendocrinology, angiogenesis, and cerebral blood flow (Voss et al., 2014). The neurobiological effects of PE are possible mediators of the benefits on cognition, particularly EF, observed in child and older adult populations (Hillman et al., 2008). Wigal and colleagues (2012) recently reviewed the

theory that PE-induced effects in youth with ADHD are due to the effects that PE has on underlying neurophysiology of ADHD.

One hypothesis suggests that PE improves cognitive functioning through increased levels of norepinephrine, dopamine, and serotonin in the prefrontal cortex (PFC), striatum, and hippocampus (Ma, 2008; Meeusen & De Meirleir, 1995; Rommel, Halperin, Mill, Asherson, & Kuntsi, 2013). This hypothesis postulates that PE-induced increases in dopamine improve working memory, attention, focus, and learning. Increases in norepinephrine enhance attention regulation, working memory, and behavioral inhibition, while increases in serotonin may further facilitate improvements in attention, affect regulation, and inhibition of hyperactive/impulsive behaviors (Salmon, 2001; Wilens & Dodson, 2004; Winter et al., 2007). Thus, the increase in dopamine and norepinephrine associated with PE is similar to the effects of stimulant medications for ADHD (Pliszka, 2005).

It has also been suggested that the cognitive effects of PE may be explained by the upregulation of BDNF (brain-derived neurotrophic factor; Knaepen, Goekint, Heyman, & Meeusen, 2010; Ma, 2008; Rommel et al., 2013; Seifert et al., 2010; Wigal et al., 2012), which is a protein involved in the survival of neurons, synapse formation, and synaptic plasticity (Reichardt, 2006). BDNF plays a critical role in the differentiation and survival of dopaminergic neurons, hippocampal functioning, and long-term potentiation for learning and memory (Voss et al., 2014). Decreased levels of central BDNF activity have been purported to play a role in the pathogenesis of ADHD; BDNF knockout mice exhibit increased hyperactivity and cognitive impairment (Tsai, 2007). A recent study with humans supported the involvement of BDNF in ADHD, as genetic variants of BDNF are linked to ADHD symptoms (Bergman, Westberg, Lichtenstein, Eriksson, & Larsson, 2011). Similar to the physiological effects of stimulant and non-stimulant medication (Fumagalli et al., 2010; Meredith, Callen, & Scheuer, 2002), PE induces upregulation of BDNF and has therefore been thought to facilitate improve-

ments in attention, inhibition, learning, memory, and mood (Hillman et al., 2008; Ma, 2008; Voss et al., 2014).

It should be noted, however, that these hypothesized neurobiological mechanisms of change are limited in that the research has largely been completed with rodents (Sontag, Tucha, Walitza, & Lange, 2010). More systematic studies of PE physiology in the context of ADHD in human samples are needed. However, as improved neuronal functioning is associated with remission of ADHD symptomatology (Bédard, Trampush, Newcorn, & Halperin, 2010; Halperin et al., 2008; Rommel et al., 2015; Shaw et al., 2013), PE has the potential of yielding enduring changes among those with ADHD (Berwid & Halperin, 2012; Halperin & Healey, 2011).

### IMPACT OF PHYSICAL EXERCISE ON ADHD

A recent systematic review identified 21 studies examining the effects of PE in children and adolescents with ADHD (for a detailed review, see Neudecker et al., 2015). Research on the effects of PE intervention programs can be classified into two groups: acute and chronic. Studies of acute effects examine efficacy in terms of the effects of PE immediately following a single bout of activity, whereas chronic studies examine efficacy in terms of the cumulative effects of PE over a specified period.

#### Acute Effects of Aerobic Physical Exercise

In their review, Neudecker and colleagues (2015) identified six studies on the acute effects of aerobic PE in children and adolescents with ADHD. Aerobic PE included either treadmill running or ergometer cycling, with variability in duration and intensity between studies. Moderate-to-large effects were reported on aspects of EF following a 20- or 30-minute treadmill exercise in two studies (Chang, Liu, Yu, & Lee, 2012; Pontifex, Saliba, Raine, Picchiotti, & Hillman, 2013). High-intensity interval treadmill running (ten 2-minute running intervals with 1 minute of rest between intervals) also was found to have small to medium effects on several aspects of attention (Medina

et al., 2010). Two studies reported that aerobic PE had small to large effects on neuroelectric measures of aspects of attention and dopaminergic activity (Pontifex et al., 2013; Tantillo, Kesick, Hynd, & Dishman, 2002), suggesting that aerobic PE may induce changes in brain activity. In the only study examining PE-induced changes in behavior, Flohr and colleagues (2004) reported significant reductions in disruptive behavior problems following 25 minutes of low to moderate ergonomic cycling. Finally, two studies using approximately similar aerobic PE duration and intensity exercises found changes in math and reading performance that were either large (Pontifex et al., 2013) or non-significant (Flohr, Saunders, Evans, & Raggi, 2004). Thus, whether aerobic PE can yield acute improvements in academic performance is unclear.

#### Chronic Effects of Regular Physical Exercise

Forming conclusions from the studies examining the chronic effects of PE in children and adolescents with ADHD is difficult due to the variability in PE regimens across studies. Studies can be categorized as either specific PE, sensorimotor PE, or mixed PE (Neudecker et al., 2015). For example, programs including tai chi would be classified as a specific PE (Hernandez-Reif, Field, & Thimas, 2001), programs focusing on perception and motor control would be classified as sensorimotor PE (Banaschewski, Bismans, Zieger, & Rothenberger, 2001; Park et al., 2013), and programs including multiple PE modalities such as aerobic and perceptual-motor water exercises would be classified as mixed PE (Chang et al., 2014).

All three categories of PE studies resulted in significant improvements on unblinded parent and teacher ratings of ADHD symptom severity in children and adolescents with ADHD (Ahmed & Mohamed, 2011; Banaschewski et al., 2001; Choi, Han, Kang, Jung, & Renshaw, 2015; Haffner, Roos, Goldstein, Parzer, & Resch, 2006; Hernandez-Reif et al., 2001; Hoza et al., 2014; Kang et al., 2011; Lufi & Parish-Plass, 2011; McKune et al., 2004), with small to large effects ( $d = .26-2.39$ ). Most PE studies reported medium to large effect sizes on objec-

tive cognitive measures aspects of EF ( $d = .37$ – $1.37$ ), including attention and inhibition, in children and adolescents with ADHD (Chang et al., 2014; Choi et al., 2014; Haffner et al., 2006; Kang, Choi, Kang, & Han, 2011; Park et al., 2013; Verret et al., 2012), although two studies failed to detect effects of yoga on attention (Jensen & Kenny, 2004) and sensorimotor training on impulse control (Banaschewski et al., 2004). Lastly, results indicated that PE interventions for children and adolescents with ADHD can improve motor skill development and physical fitness, as evidenced by PE having moderate to large effects ( $d = .31$ – $2.82$ ) on measures of motor performance (Ahmed & Mohamed, 2011; Banaschewski et al., 2001; Chang et al., 2014; McKune et al., 2004; Pan et al., 2015; Verret et al., 2012).

### Summary of the Impacts of PE on ADHD

Overall, the existing literature provides preliminary support for acute and chronic effects of PE on ADHD symptomatology and common areas of impairments in children and adolescents. In addition, PE has been associated with improvements on physiological measures associated with ADHD. However, there are no known studies demonstrating that improvements on physiological measures mediate improvements in ADHD symptomatology or impairments. Thus, it is unclear if observed improvements in impairment and symptoms are the result of changes in neurobiological factors. Most of these studies suffer from one or more limitations, such as limited sample size, subjective measures, nonrandom assignment, or absent or inadequate comparison groups.

### PHYSICAL EXERCISE AS A POSSIBLE INTERVENTION FOR EMERGING ADULTS WITH ADHD

Although preliminary findings suggest that PE can have beneficial effects on the cognitive and behavioral functioning of children and adolescents with ADHD, it remains unknown whether the effectiveness of PE as a therapeutic tool varies across the developmental spectrum. Although it may be suspected that PE would yield the greatest effects dur-

ing the developmental period in which the brain is most malleable, this notion has not been empirically tested. However, the PFC is among the latest neural structures to mature, continuing to develop throughout the mid-20s (Giedd, 2004), and increases in BDNF have been shown to be correlated with sustained attention among adults (Halperin & Healey, 2011). Evidence of the beneficial effects of PE on cognition in elderly adults (e.g., Angevaren et al., 2008) reinforces the possibility that PE could be beneficial for ameliorating core symptoms and impairments of ADHD beyond childhood.

While PE appears to be a plausibly *efficacious* intervention for ADHD in emerging adulthood, important consideration should be given to the degree to which it may be *effective*. That is, PE interventions may be efficacious in tightly controlled, ideal circumstances, but may not be effective in real world clinical settings. For those with ADHD, the transition to emerging adulthood is associated with almost complete treatment disengagement (McCarthy et al., 2009). Similarly, adherence to a regimen of consistent PE is notoriously poor over time (e.g., Garber et al., 2011). Unlike school-aged children who can have PE interventions incorporated into school schedules (Hoza et al., 2014) or are mandated by their parents to receive treatment, emerging adults must voluntarily engage, allocate time, and typically pay for treatments. Thus, it is important to consider the extent to which PE interventions can be effective outside of school settings. The following section outlines the research on PE interventions among adults with major depressive disorder (MDD) to inform the possible effectiveness of PE as an intervention tool for those with ADHD in emerging adulthood.

### EFFECTIVENESS OF PHYSICAL EXERCISE INTERVENTIONS FOR MAJOR DEPRESSIVE DISORDER

The use of exercise in improving mental health is a growing research area that has documented positive effects across a variety of domains, including depression, anxiety, and stress (Salmon, 2001), as well as neurodegenerative diseases (Deslandes et al., 2010). Researchers cite

the similar neurobiological mechanisms underlying the positive effects of PE for ADHD and MDD. This includes PE-induced increases in the synthesis and release of neurotransmitters and BDNF, which are thought to result in neurogenesis, angiogenesis, and neuroplasticity (Portugal et al., 2013). The mental disorder which has received the most research in PE is MDD. Theories underlying the pathophysiology of MDD often involve reductions of norepinephrine and serotonin (Lopez-Munoz & Alamo, 2009) and hypothalamic-pituitary-adrenal (HPA) axis hyperactivity due to increased levels of cortisol and corticotropin-releasing factor (Palazidou, 2012).

In follow-up studies, open trials, and randomized control trials, symptom reduction of MDD has consistently been correlated with regular PE (e.g., Blumenthal et al., 2007; Deslandes et al., 2010; Dunn, Trivedi, Kampert, Clark, & Chambliss, 2005; Singh et al., 2005). Both strength and aerobic PE evidence moderate improvements in MDD symptoms (standardized mean difference [SMD] = .61), with no significant differences between the two types of interventions (Silveira et al., 2013). Silveira and colleagues (2013) noted that a combination of moderate-intensity aerobic training and high-intensity strength training may provide more positive benefits than other PE regimens. While PE appears to be efficacious in reducing MDD symptom severity, the impact of PE is dependent upon level of adherence to a PE regimen (Portugal et al., 2013).

As alluded to previously, the positive mental health and cognitive effects associated with PE are predicated on regular engagement in PE (Garber et al., 2011), with high rates of *inactivity* attenuating beneficial effects (Guthold, Ono, Strong, Chatterji, & Morabia, 2008; Vayman & Gomez-Pinilla, 2006; Voss et al., 2014). The affective response to PE has been shown to predict the level of engagement 6 and 12 months after a PE session (Williams et al., 2008). In the case of aerobic PE, Ekkekakis and Petruzzello (1999) suggested that the optimal intensity to provoke the most positive affective response is moderate to high and is near ventilatory threshold—the point at which oxygen delivery to muscles

becomes a limiting factor (~65%  $VO_{2max}$ ; Caiozzo et al., 1982). Aerobic regimens that tend to elicit positive affect include moderate aerobic PE that is below the ventilatory threshold (Reed & Ones, 2006). Although aerobic PE above the ventilatory threshold may be perceived as threatening and generate negative affective states (Ekkekakis, 2003), exercise regimens involving intervals of high-intensity aerobic PE elicit greater pleasure than continuous, moderate aerobic PE (Bartlett et al., 2011). In the case of strength training, little research has examined the optimal intensity to provoke the most positive affective response, with studies comparing low- and high-intensity strength training yielding inconsistent results (Portugal et al., 2013). Thus, there appears to be no consensus regarding the optimal strength training intensity on mood.

#### CLINICAL RECOMMENDATIONS FOR ADHD

Despite the absence of research documenting the effectiveness of PE in emerging adults with ADHD, there are numerous reasons for mental health providers to recommend PE as an adjunct to psychosocial and/or pharmacological treatments for clients of all ages. First, the existing literature in children and adolescents with ADHD provides evidence suggesting PE may yield improvements in cognitive and behavioral functioning in emerging adults. Second, regardless of ADHD status, PE is recommended for all adults to improve overall health and reduce the likelihood of future diseases (U.S. Department of Health and Human Services, 2008). However, PE is arguably more beneficial for emerging adults with ADHD due to their increased risk for poor health outcomes (Barkley et al., 2008; Nigg, 2013). Third, PE ameliorates depressive and anxiety symptoms (De Moor, Beem, Stubbe, Boomsma, & De Geus, 2006), both of which are commonly comorbid in emerging adults with ADHD (Meinzer et al., 2013; Milberger, Biederman, Faraone, Murphy, & Tsuang, 1995). Fourth, emerging adults with ADHD are 2 to 5 times more likely to experience sleeping problems (Gau et al., 2007), which may be improved by

regular PE (Driver & Taylor, 2000). Although the plausibility of PE improving sleep patterns with ADHD samples has not been empirically tested, it is both relevant and important because a common side effect of stimulant medication is difficulty sleeping, which is negatively associated with mood regulation, academic performance, and domains of EF, including sustained attention, working memory, and inhibition (Danner & Phillips, 2008; Durmer & Dinges, 2005; Fafrowicz et al., 2010; Van Dongen et al., 2003). Lastly, for emerging adults without PE medical contraindications, safe levels of PE are a “do no harm” intervention tool with no known persistent adverse effects (Hoza & Smith, 2015, p. 4). In a similar vein, Arnold, Hurt, and Lofthouse (2013) have argued that some treatments are “safe, easy, cheap, and sensible (SECS) enough to be tried while awaiting better research” (p. 381). Thus, in spite of the limited evidence base, we are advising clinicians to consider including moderate-to-vigorous physical exercise as a recommendation for all clients with ADHD who have been approved for exercise by their physician.

#### FUTURE RESEARCH DIRECTIONS

Given that research on interventions for ADHD are scarce for emerging adults who often struggle during this developmental period (Fleming & McMahon, 2012; LaCount, Hartung, Canu, & Knouse, 2018) and that adjustment during this period tends to predict adaptive functioning in adulthood (Schulenberg, Sameroff, & Cicchetti, 2004), research exploring the potential benefits of PE in this population is warranted. Specifically, research is needed to test our aforementioned hypotheses regarding the possible effectiveness and efficacy of PE interventions for this age group. Relevant to PE interventions for ADHD among school-aged children and emerging adults, several critical questions remain:

1. What is the optimal dosage and modality of physical exercise and/or physical activity?
2. What is the minimum necessary intensity and duration needed to achieve a measurable and clinically

significant effect (e.g., 30% symptom reduction; Zylowska et al., 2008)?

3. Are observed PE effects moderated by individual characteristics? In addition to the large variability in PE protocols, there was considerable heterogeneity between samples with a lack of consideration for important factors (e.g., medication status, comorbidities).
4. Will the impact of PE on cognitive function on laboratory and questionnaire measures generalize to real world benefits? For example, although cognitive training interventions for ADHD often report improvements in cognitive function and ADHD symptomatology, the improvements often fail to have ecological validity (Cortese et al., 2015).
5. Do physiological mechanisms mediate positive effects of PE on ADHD-related outcomes? In this respect, it is also important to consider other possible explanations for PE-induced improvements. It is often overlooked that many forms of PE require cognitive engagement (e.g., group games) that might explain improvements in cognitive function (Best, 2010). For example, it seems plausible that the mechanisms underlying the positive effects of PE may be similar to those of psychological mechanisms associated with mindfulness meditation training. Mindfulness meditation has been described as a psychological process that includes purposefully orienting attention to the present moment and approaching experiences in the present with curiosity, openness, and acceptance (Bishop et al., 2004). Although not empirically tested, it is intuitively plausible that, for example, engaging in moderate-intensity PE induces present-mindedness. Indeed, PE interventions and mindfulness meditation interventions cite a similar neurobiological mechanism (e.g., anterior cingulate cortex) to explain improvements in cognitive and behavioral outcomes among individuals with ADHD

(Mitchell, Zylowska, & Kollins, 2015).

6. What is the size of the effect and incremental benefit of PE relative to established ADHD treatments? Although existing studies have compared PE interventions against other interventions and included medicated and unmedicated samples, the individual and additive benefit of PE interventions to evidenced-based treatments for ADHD has not been systematically studied.

Patrick A. LaCount, M.S., is a doctoral candidate in the Department of Psychology at the University of Wyoming and a predoctoral intern at Duke University Medical Center. Cynthia M. Hartung, Ph.D., is Associate Professor and Director of the Psychology Clinic in the Psychology Department at the University of Wyoming. Dr. Hartung can be contacted at [chartung@uwyo.edu](mailto:chartung@uwyo.edu). This project was supported in part by the Charles Ksir Summer Research Fellowship and an Institutional Development Award (IDeA) from the National Institute of General Medical Sciences of the National Institutes of Health under grant number 2P20GM103432. Its contents are solely the responsibility of the authors and do not represent the official views of NIH or Charles Ksir. We thank Derek Smith, Joshua Clapp, and Christine McKibbin for reviewing earlier versions of this manuscript.

## REFERENCES

Åberg, M. I., Pedersen, N. L., Torén, K., Svartengren, M., Bäckstrand, B., Johnsson, T., ... Kuhn, H. G. (2009). Cardiovascular fitness is associated with cognition in young adulthood. *PNAS Proceedings of the National Academy of Sciences of The United States of America*, *106*, 20906-20911.

Ahmed, G., & Mohamed, S. (2011). Effect of regular aerobic exercises on behavioral, cognitive and psychological response in patients with attention deficit-hyperactivity disorder. *Life Science Journal*, *8*, 366-371.

American Psychiatric Association (APA). (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: Author.

Angevaren, M., Aufdemkampe, G., Verhaar, H. J., Aleman, A., & Vanhees, L. (2008). Physical activity and enhanced fitness to improve cognitive function in older people without known cognitive impairment. *Co-*

*chrane Database Systematic Reviews*, *3*, CD005381.

Arnett, J. J. (2000). Emerging adulthood: A theory of development from the late teens through the twenties. *American Psychologist*, *55*, 469-480.

Arnold, L. E., Hurt, E., & Lofthouse, N. (2013). Attention-deficit/hyperactivity disorder: Dietary and nutritional treatments. *Child and Adolescent Psychiatry Clinics of North America*, *22*, 381-402.

Arnsten, A. F. T. (2000). Genetics of childhood disorders: XVIII. ADHD, Part 2: Norepinephrine has a critical modulatory influence on prefrontal cortical function. *Journal of the American Academy of Child & Adolescent Psychiatry*, *39*, 1201-1203.

Arnsten, A. T. (2009). Toward a new understanding of attention-deficit hyperactivity disorder pathophysiology: An important role for prefrontal cortex dysfunction. *CNS Drugs*, *23*(Suppl. 1), 33-41.

Banaschewski, T., Bismans, F., Zieger, H., & Rothenberger, A. (2001). Evaluation of sensorimotor training in children with ADHD. *Perceptual and Motor Skills*, *92*(1), 137-149.

Barkley, R. A. (1997). Behavioral inhibition, sustained attention, and executive functions: Constructing a unifying theory of ADHD. *Psychological Bulletin*, *121*, 65-94.

Barkley, R. A., Murphy, K. R., & Fischer, M. (2008). *ADHD in adults: What the science says*. New York, NY: Guilford.

Bartlett, J. D., Close, G. L., MacLaren, D. P., Gregson, W., Drust, B., & Morton, J. P. (2011). High-intensity interval running is perceived to be more enjoyable than moderate-intensity continuous exercise: Implications for exercise adherence. *Journal of Sports Sciences*, *29*, 547-553.

Bédard, A. C. V., Trampush, J. W., Newcorn, J. H., & Halperin, J. M. (2010). Perceptual and motor inhibition in adolescents/young adults with childhood-diagnosed ADHD. *Neuropsychology*, *24*, 424.

Bergman, O., Westberg, L., Lichtenstein, P., Eriksson, E., & Larsson, H. (2011). Study on the possible association of brain-derived neurotrophic factor polymorphism with the developmental course of symptoms of attention deficit and hyperactivity. *International Journal of Neuropsychopharmacology*, *14*, 1367-1376.

Berridge, C. W., & Devilbiss, D. M. (2011). Psychostimulants as cognitive enhancers: The prefrontal cortex, catecholamines, and attention-deficit/hyperactivity disorder. *Biological Psychiatry*, *69*, e101-e111.

Berwid, O. G., & Halperin, J. M. (2012). Emerging support for a role of exercise in attention-deficit/hyperactivity disorder

intervention planning. *Current Psychiatry Reports*, *14*, 543-551.

Best, J. R. (2010). Effects of physical activity on children's executive function: Contributions of experimental research on aerobic exercise. *Developmental Review*, *30*, 331-351.

Bishop, S. R., Lau, M., Shapiro, S., Carlson, L., Anderson, N. D., Carmody, J., ... Devins, G. (2004). Mindfulness: A proposed operational definition. *Clinical Psychology: Science and Practice*, *11*, 230-241.

Blumenthal, J. A., Babyak, M. A., Doraiswamy, P. M., Watkins, L., Hoffman, B. M., Barbour, K. A., ... Sherwood, A. (2007). Exercise and pharmacotherapy in the treatment of major depressive disorder. *Psychosomatic Medicine*, *69*, 587.

Caiozzo, V. J., Davis, J. A., Ellis, J. F., Azus, J. L., Vandagriff, R., Prietto, C. A., & McMaster, W. C. (1982). A comparison of gas exchange indices used to detect the anaerobic threshold. *Journal of Applied Physiology*, *53*, 1184-1189.

Caspersen, C. J., Powell, K. E., & Christenson, G. M. (1985). Physical activity, exercise, and physical fitness: Definitions and distinctions for health-related research. *Public Health Reports*, *100*, 126.

Castellanos, F. X., & Tannock, R. (2002). Neuroscience of attention-deficit/hyperactivity disorder: The search for endophenotypes. *Nature Reviews Neuroscience*, *3*, 617-628.

Chang, Y., Hung, C., Huang, C., Hatfield, B. D., & Hung, T. (2014). Effects of an aquatic exercise program on inhibitory control in children with ADHD: A preliminary study. *Archives of Clinical Neuropsychology*, *29*, 217-223.

Chang, Y. K., Liu, S., Yu, H. H., & Lee, Y. H. (2012). Effect of acute exercise on executive function in children with attention deficit hyperactivity disorder. *Archives of Clinical Neuropsychology*, *27*, 225-237.

Choi, J. W., Han, D. H., Kang, K. D., Jung, H. Y., & Renshaw, P. F. (2015). Aerobic exercise and attention deficit hyperactivity disorder: Brain research. *Medicine and Science in Sports and Exercise*, *47*, 33-39.

Cortese, S., Ferrin, M., Brandeis, D., Buitelaar, J., Daley, D., Dittmann, R. W., ... European ADHD Guidelines Group. (2015). Cognitive training for attention-deficit/hyperactivity disorder: Meta-analysis of clinical and neuropsychological outcomes from randomized controlled trials. *Journal of the American Academy of Child & Adolescent Psychiatry*, *54*, 164-174.

Danner, F., & Phillips, B. (2008). Adolescent sleep, school start times, and teen motor vehicle crashes. *Journal of Clinical Sleep Medicine*, *4*, 533-535.



- Davis, C. L., Tomporowski, P. D., McDowell, J. E., Austin, B. P., Miller, P. H., Yanasak, N. E., & ... Naglieri, J. A. (2011). Exercise improves executive function and achievement and alters brain activation in overweight children: A randomized, controlled trial. *Health Psychology, 30*, 91-98.
- De Moor, M. H. M., Beem, A. L., Stubbe, J. H., Boomsma, D. I., & De Geus, E. J. C. (2006). Regular exercise, anxiety, depression and personality: A population-based study. *Preventive Medicine, 42*, 273-279.
- Deslandes, A. C., Moraes, H., Alves, H., Pompeu, F. A. M. S., Silveira, H., Mouta, R., ... Coutinho, E. S. F. (2010). Effect of aerobic training on EEG alpha asymmetry and depressive symptoms in the elderly: A 1-year follow-up study. *Brazilian Journal of Medical and Biological Research, 43*, 585-592.
- Dickstein, S. G., Bannon, K., Castellanos, F. X., & Milham, M. P. (2006). The neural correlates of attention deficit hyperactivity disorder: An ALE meta-analysis. *Journal of Child Psychology and Psychiatry, 47*, 1051-1062.
- Driver, H. S., & Taylor, S. R. (2000). Exercise and sleep. *Sleep Medicine Reviews, 4*, 387-402.
- Dunn, A. L., Trivedi, M. H., Kampert, J. B., Clark, C. G., & Chambliss, H. O. (2005). Exercise treatment for depression: Efficacy and dose response. *American Journal of Preventive Medicine, 28*, 1-8.
- Durmer, J. S., & Dinges, D. F. (2005). Neurocognitive consequences of sleep deprivation. *Seminars in Neurology, 25*, 117-129.
- Durstewitz, D., Kelc, M., & Güntürkün, O. (1999). A neurocomputational theory of the dopaminergic modulation of working memory functions. *Journal of Neuroscience, 19*, 2807-2822.
- Durston, S. (2003). A review of the biological bases of ADHD: What have we learned from imaging studies? *Mental Retardation and Developmental Disabilities Research Reviews, 9*, 184-195.
- Durston, S., & Konrad, K. (2007). Integrating genetic, psychopharmacological and neuroimaging studies: A converging methods approach to understanding the neurobiology of ADHD. *Developmental Review, 27*, 374-395.
- Ekkekakis, P. (2003). Pleasure and displeasure from the body: Perspectives from exercise. *Cognition & Emotion, 17*, 213-239.
- Ekkekakis, P., & Petruzzello, S. J. (1999). Acute aerobic exercise and affect: Current status, problems and prospects regarding dose-response. *Sports Medicine, 5*, 337-374.
- Etnier, J. L., Nowell, P. M., Landers, D. M., & Sibley, B. A. (2006). A meta-regression to examine the relationship between aerobic fitness and cognitive performance. *Brain Research Reviews, 52*, 119-130.
- Fafrowicz, M., Oginska, H., Mojsa-Kaja, J., Marek, T., Golonka, K., & Tucholska, K. (2010). Chronic sleep deficit and performance of a sustained attention task: An electrooculography study. *Chronobiology International, 27*, 934-944.
- Fleming, A. P., & McMahon, R. J. (2012). Developmental context and treatment principles for ADHD among college students. *Clinical Child and Family Psychology Review, 15*, 303-329.
- Flohr, J. A., Saunders, M. J., Evans, S. W., & Raggi, V. (2004). Effects of physical activity on academic performance and behavior in children with ADHD. *Medicine & Science in Sports & Exercise, 36*, S145-S146.
- Fumagalli, F., Cattaneo, A., Caffino, L., Ibba, M., Racagni, G., Carboni, E., ... Riva, M. A. (2010). Sub-chronic exposure to atomoxetine up-regulates BDNF expression and signalling in the brain of adolescent spontaneously hypertensive rats: Comparison with methylphenidate. *Pharmacological Research, 62*, 523-529.
- Gapin, J. I., Labban, J. D., & Etnier, J. L. (2011). The effects of physical activity on attention deficit hyperactivity disorder symptoms: The evidence. *Preventive Medicine, 52*, S70-S74.
- Garber, C. E., Blissmer, B., Deschenes, M. R., Franklin, B. A., Lamonte, M. J., Lee, I. M., ... Swain, D. P. (2011). American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: Guidance for prescribing exercise. *Medicine & Science in Sports & Exercise, 43*, 1334-1359.
- Gau, S. F., Kessler, R. C., Tseng, W., Wu, Y., Chiu, Y., Yeh, C., & Hwu, H. (2007). Association between sleep problems and symptoms of attention-deficit/hyperactivity disorder in young adults. *Sleep: Journal of Sleep and Sleep Disorders Research, 30*, 195-201.
- Giedd, J. N. (2004). Structural magnetic resonance imaging of the adolescent brain. *Annals of the New York Academy of Sciences, 1021*, 77-85.
- Guthold, R., Ono, T., Strong, K. L., Chatterji, S., & Morabia, A. (2008). Worldwide variability in physical inactivity: A 51-country survey. *American Journal of Preventive Medicine, 34*, 486-494.
- Haffner, J., Roos, J., Goldstein, N., Parzer, P., & Resch, F. (2006). The effectiveness of body-oriented methods of therapy in the treatment of attention-deficit hyperactivity disorder (ADHD): Results of a controlled pilot study. *Zeitschrift für Kinder- und Jugendpsychiatrie und Psychotherapie, 34*, 37-47.
- Halperin, J. M., & Healey, D. M. (2011). The influences of environmental enrichment, cognitive enhancement, and physical exercise on brain development: Can we alter the developmental trajectory of ADHD? *Neuroscience & Biobehavioral Reviews, 35*, 621-634.
- Halperin, J. M., Trampush, J. W., Miller, C. J., Marks, D. J., & Newcorn, J. H. (2008). Neuropsychological outcome in adolescents/young adults with childhood ADHD: Profiles of persisters, remitters and controls. *Journal of Child Psychology and Psychiatry, 49*, 958-966.
- Hernandez-Reif, M., Field, T. M., & Thimas, E. (2001). Attention deficit hyperactivity disorder: Benefits from tai chi. *Journal of Bodywork and Movement Therapies, 5*, 120-123.
- Hervey, A. S., Epstein, J. N., & Curry, J. F. (2004). Neuropsychology of adults with attention-deficit/hyperactivity disorder: A meta-analytic review. *Neuropsychology, 18*, 485-503.
- Hillman, C. H., Erickson, K. I., & Kramer, A. F. (2008). Be smart, exercise your heart: Exercise effects on brain and cognition. *Nature Reviews Neuroscience, 9*, 58-65.
- Hoza, B., & Smith, A. L. (2015). Is aerobic physical activity a viable management strategy for ADHD? *The ADHD Report, 23*, 1-5.
- Hoza, B., Smith, A. L., Shoulberg, E. K., Linnea, K. S., Dorsch, T. E., Blazo, J. A., ... McCabe, G. P. (2014). A randomized trial examining the effects of aerobic physical activity on attention-deficit/hyperactivity disorder symptoms in young children. *Journal of Abnormal Child Psychology, 43*, 655-667.
- Jensen, P. S., & Kenny, D. T. (2004). The effects of yoga on the attention and behavior of boys with attention-deficit/hyperactivity disorder (ADHD). *Journal of Attention Disorders, 7*, 205-216.
- Kang, K. D., Choi, J. W., Kang, S. G., & Han, D. H. (2011). Sports therapy for attention, cognitions and sociality. *International Journal of Sports Medicine, 32*, 953-959.
- Knaepen, K., Goekint, M., Heyman, E. M., & Meeusen, R. (2010). Neuroplasticity: Exercise-induced response of peripheral brain-derived neurotrophic factor. *Sports Medicine, 40*, 765-801.
- Knight, L. A., Rooney, M., & Chronis-Tuscano, A. (2008). Psychosocial treatments for attention-deficit/hyperactivity disorder. *Current Psychiatry Reports, 10*, 412-418.
- Kramer, A. F., & Erickson, K. I. (2007). Capitalizing on cortical plasticity: Influence of physical activity on cognition and brain function. *Trends in Cognitive Sciences, 11*, 342-348.
- LaCount, P. A., Hartung, C. M., Canu, W. H., & Knouse, L. E. (2018). Interventions for transitioning adolescents with ADHD to emerging adulthood: Developmental context and empirically supported treatment principles. Manuscript under review.

- Lopez-Munoz, F., & Alamo, C. (2009). Monoaminergic neurotransmission: The history of the discovery of antidepressants from 1950s until today. *Current Pharmaceutical Design*, 15, 1563-1586.
- Loprinzi, P. D., Herod, S. M., Cardinal, B. J., & Noakes, T. D. (2013). Physical activity and the brain: A review of this dynamic, bi-directional relationship. *Brain Research*, 1539, 95-104.
- Lufi, D., & Parish-Plass, J. (2011). Sport-based group therapy program for boys with ADHD or with other behavioral disorders. *Child & Family Behavior Therapy*, 33, 217-230.
- Luman, M., Oosterlaan, J., & Sergeant, J. A. (2005). The impact of reinforcement contingencies on AD/HD: A review and theoretical appraisal. *Clinical Psychology Review*, 25, 183-213.
- Ma, Q. (2008). Beneficial effects of moderate voluntary physical exercise and its biological mechanisms on brain health. *Neuroscience Bulletin*, 24, 265-270.
- Makris, N., Biederman, J., Valera, E. M., Bush, G., Kaiser, J., Kennedy, D. N., ... Seidman, L. J. (2007). Cortical thinning of the attention and executive function networks in adults with attention-deficit/hyperactivity disorder. *Cerebral Cortex*, 17, 1364-1375.
- McCarthy, S., Asherson, P., Coghill, D., Hollis, C., Murray, M., Potts, L., ... Wong, I. C. (2009). Attention-deficit hyperactivity disorder: Treatment discontinuation in adolescents and young adults. *British Journal of Psychiatry*, 194, 273-277.
- McKune, A. J., Pautz, J., & Lomjard, J. (2004). Behavioural response to exercise in children with attention-deficit/hyperactivity disorder. *South African Journal of Sports Medicine*, 15, 17-21.
- Medina, J. A., Netto, T. L., Muszkat, M., Medina, A. C., Botter, D., Orbetelli, R., ... Miranda, M. C. (2010). Exercise impact on sustained attention of ADHD children, methylphenidate effects. *Attention Deficit and Hyperactivity Disorders*, 2, 49-58.
- Meeusen, R., & De Meirleir, K. (1995). Exercise and brain neurotransmission. *Sports Medicine*, 20, 160-188.
- Meinzer, M. C., Lewinsohn, P. M., Pettit, J. W., Seeley, J. R., Gau, J. M., Chronis-Tuscano, A., & Waxmonsky, J. G. (2013). Attention-deficit/hyperactivity disorder in adolescence predicts onset of major depressive disorder through early adulthood. *Depression and Anxiety*, 30, 546-553.
- Meredith, G. E., Callen, S., & Scheuer, D. A. (2002). Brain-derived neurotrophic factor expression is increased in the rat amygdala, piriform cortex and hypothalamus following repeated amphetamine administration. *Brain Research*, 949, 218-227.
- Milberger, S., Biederman, J., Faraone, S. V., Murphy, J., & Tsuang, M. T. (1995). Attention deficit hyperactivity disorder and comorbid disorders: Issues of overlapping symptoms. *American Journal of Psychiatry*, 152, 1793-1799.
- Mitchell, J. T., Zylowska, L., & Kollins, S. H. (2015). Mindfulness meditation training for attention-deficit/hyperactivity disorder in adulthood: Current empirical support, treatment overview, and future directions. *Cognitive and Behavioral Practice*, 22, 172-191.
- Molina, B. G., Hinshaw, S. P., Swanson, J. M., Arnold, L. E., Vitiello, B., Jensen, P. S., ... Houck, P. R. (2009). The MTA at 8 years: Prospective follow-up of children treated for combined-type ADHD in a multisite study. *Journal of the American Academy of Child & Adolescent Psychiatry*, 48, 484-500.
- Neudecker, C., Mewes, N., Reimers, A. K., & Woll, A. (2015). Exercise intervention in children and adolescents with ADHD: A systematic review. *Journal of Attention Disorders*. Ahead of print.
- Nigg, J. T. (2013). Attention-deficit/hyperactivity disorder and adverse health outcomes. *Clinical Psychology Review*, 33, 215-228.
- Nigg, J. T., Hinshaw, S. P., Carte, E. T., & Treuting, J. J. (1998). Neuropsychological correlates of childhood attention-deficit/hyperactivity disorder: Explainable by comorbid disruptive behavior or reading problems? *Journal of Abnormal Psychology*, 107, 468-480.
- Nigg, J. T., Stavro, G., Ettenhofer, M., Hambrick, D. Z., Miller, T., & Henderson, J. M. (2005). Executive functions and ADHD in adults: Evidence for selective effects on ADHD symptom domains. *Journal of Abnormal Psychology*, 114, 706-717.
- Palazidou, E. (2012). The neurobiology of depression. *British Medical Bulletin*, 101, 127-145.
- Pan, C. Y., Tsai, C. L., Chu, C. H., Sung, M. C., Huang, C. Y., & Ma, W. Y. (2015). Effects of physical exercise intervention on motor skills and executive functions in children with ADHD: A pilot study. *Journal of Attention Disorders*. Ahead of print.
- Park, M. S., Byun, K. W., Park, Y. K., Kim, M. H., Jung, S. H., & Kim, H. (2013). Effect of complex treatment using visual and auditory stimuli on the symptoms of attention deficit/hyperactivity disorder in children. *Journal of Exercise Rehabilitation*, 9, 316-325.
- Pennington, B. F. (2006). From single to multiple deficit models of developmental disorders. *Cognition*, 101, 385-413.
- Piepmeier, A. T., Shih, C. H., Whedon, M., Williams, L. M., Davis, M. E., Henning, D. A., ... Etnier, J. L. (2015). The effect of acute exercise on cognitive performance in children with and without ADHD. *Journal of Sport and Health Science*, 4, 97-104.
- Pliszka, S. R. (2005). The neuropsychopharmacology of attention-deficit/hyperactivity disorder. *Biological Psychiatry*, 57, 1385-1390.
- Pontifex, M. B., Saliba, B. J., Raine, L. B., Picchietti, D. L., & Hillman, C. H. (2013). Exercise improves behavioral, neurocognitive, and scholastic performance in children with attention-deficit/hyperactivity disorder. *Journal of Pediatrics*, 162, 543-551.
- Portugal, E. M., Cevada, T., Monteiro-Junior, R. S., Guimaraes, T. T., da Cruz Rubini, E., Lattari, E., ... Deslandes, A. C. (2013). Neuroscience of exercise: From neurobiology mechanisms to mental health. *Neuro-psychobiology*, 68, 1-14.
- Reed, J., & Ones, D. S. (2006). The effect of acute aerobic exercise on positive activated affect: A meta-analysis. *Psychology of Sport and Exercise*, 7, 477-514.
- Reichardt, L. F. (2006). Neurotrophin-regulated signalling pathways. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, 361, 1545-1564.
- Rommel, A. S., Halperin, J. M., Mill, J., Asherson, P., & Kuntsi, J. (2013). Protection from genetic diathesis in attention-deficit/hyperactivity disorder: Possible complementary roles of exercise. *Journal of the American Academy of Child & Adolescent Psychiatry*, 52, 900-910.
- Rommel, A. S., Lichtenstein, P., Rydell, M., Kuja-Halkola, R., Asherson, P., Kuntsi, J., & Larsson, H. (2015). Is physical activity causally associated with symptoms of attention-deficit/hyperactivity disorder? *Journal of the American Academy of Child & Adolescent Psychiatry*, 54, 555-570.
- Salmon, P. (2001). Effects of physical exercise on anxiety, depression, and sensitivity to stress: A unifying theory. *Clinical Psychology Review*, 21, 33-61.
- Scheres, A., Milham, M. P., Knutson, B., & Castellanos, F. X. (2007). Ventral striatal hypo-responsiveness during reward anticipation in attention-deficit/hyperactivity disorder. *Biological Psychiatry*, 61, 720-724.
- Schoechlin, C., & Engel, R. R. (2005). Neuropsychological performance in adult attention-deficit hyperactivity disorder: Meta-analysis of empirical data. *Archives of Clinical Neuropsychology*, 20, 727-744.
- Schulenberg, J. E., Sameroff, A. J., & Cicchetti, D. (2004). The transition to adulthood as a critical juncture in the course of psychopathology and mental health. *Development and Psychopathology*, 16, 799-806.
- Seifert, T., Brassard, P., Wissenberg, M., Rasmussen, P., Nordby, P., Stallknecht, B., ... Secher, N. H. (2010). Endurance training enhances BDNF release from the human

- brain. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 298, R372-R377.
- Shaw, P., Malek, M., Watson, B., Greenstein, D., de Rossi, P., & Sharp, W. (2013). Trajectories of cerebral cortical development in childhood and adolescence and adult attention-deficit/hyperactivity disorder. *Biological Psychiatry*, 74, 599-606.
- Sibley, B. A., & Beilock, S. L. (2007). Exercise and working memory: An individual differences investigation. *Journal of Sport & Exercise Psychology*, 29, 783-791.
- Sibley, B. A., & Etnier, J. L. (2003). The relationship between physical activity and cognition in children: A meta-analysis. *Pediatric Exercise Science*, 15, 243-256.
- Silveira, H., Moraes, H., Oliveira, N., Coutinho, E. S. F., Laks, J., & Deslandes, A. (2013). Physical exercise and clinically depressed patients: A systematic review and meta-analysis. *Neuropsychobiology*, 67, 61-68.
- Singh, N. A., Stavrinou, T. M., Scarbek, Y., Galambos, G., Liber, C., & Singh, M. A. F. (2005). A randomized controlled trial of high versus low intensity weight training versus general practitioner care for clinical depression in older adults. *The Journals of Gerontology, Series A: Biological Sciences and Medical Sciences*, 60, 768-776.
- Solanto, M. V. (2002). Dopamine dysfunction in AD/HD: Integrating clinical and basic neuroscience research. *Behavioural Brain Research*, 130(1-2), 65-71.
- Sontag, T. A., Tucha, O., Walitza, S., & Lange, K. W. (2010). Animal models of attention deficit/hyperactivity disorder (ADHD): A critical review. *Attention Deficit and Hyperactivity Disorders*, 2, 1-20.
- Sonuga-Barke, E. J. (2002). Psychological heterogeneity in AD/HD: A dual pathway model of behaviour and cognition. *Behavioural Brain Research*, 130, 29-36.
- Spencer, T., Biederman, J., Wilens, T., Harding, M., O'Donnell, D., & Griffin, S. (1996). Pharmacotherapy of attention-deficit hyperactivity disorder across the life cycle. *Journal of the American Academy of Child & Adolescent Psychiatry*, 35, 409-432.
- Ströhle, A., Stoy, M., Wrase, J., Schwarzer, S., Schlagenhauf, F., Huss, M., ... Heinz, A. (2008). Reward anticipation and outcomes in adult males with attention-deficit/hyperactivity disorder. *NeuroImage*, 39, 966-972.
- Tantillo, M., Kesick, C. M., Hynd, G. W., & Dishman, R. K. (2002). The effects of exercise on children with attention-deficit hyperactivity disorder. *Medicine & Science in Sports & Exercise*, 1, 203-212.
- Taylor, A. F., & Kuo, F. E. (2009). Children with attention deficits concentrate better after walk in the park. *Journal of Attention Disorders*, 12, 402-409.
- Tomprowski, P. D., Lambourne, K., & Okumura, M. S. (2011). Physical activity interventions and children's mental function: An introduction and overview. *Preventive Medicine: An International Journal Devoted to Practice and Theory*, 52(Suppl.), S3-S9.
- Trost, S. G., Owen, N., Bauman, A. E., Sallis, J. F., & Brown, W. (2002). Correlates of adults' participation in physical activity: Review and update. *Medicine & Science in Sports & Exercise*, 34, 1996-2001.
- Tsai, S. J. (2007). Attention-deficit hyperactivity disorder may be associated with decreased central brain-derived neurotrophic factor activity: Clinical and therapeutic implications. *Medical Hypotheses*, 68, 896-899.
- U.S. Department of Health and Human Services. (2008). *2008 physical activity guidelines for Americans*. <http://www.health.gov/paguidelines/>
- Van Dongen, H. A., Maislin, G., Mullington, J. M., & Dinges, D. F. (2003). The cumulative cost of additional wakefulness: Dose-response effects on neurobehavioral functions and sleep physiology from chronic sleep restriction and total sleep deprivation. *Sleep: Journal of Sleep and Sleep Disorders Research*, 26, 117-126.
- Vaynman, S., & Gomez-Pinilla, F. (2006). Revenge of the "sit": How lifestyle impacts neuronal and cognitive health through molecular systems that interface energy metabolism with neuronal plasticity. *Journal of Neuroscience Research*, 84, 699-715.
- Verret, C., Guay, M., Berthiaume, C., Gardiner, P., & Béliveau, L. (2012). A physical activity program improves behavior and cognitive functions in children with ADHD: An exploratory study. *Journal of Attention Disorders*, 16, 71-80.
- Voss, M. W., Carr, L. J., Clark, R., & Weng, T. (2014). Revenge of the "sit" II: Does lifestyle impact neuronal and cognitive health through distinct mechanisms associated with sedentary behavior and physical activity? *Mental Health and Physical Activity*, 7, 9-24.
- Wigal, S. B., Emmerson, N., Gehricke, J., & Galassetti, P. (2012). Exercise: Applications to childhood ADHD. *Journal of Attention Disorders*, 17, 279-290.
- Wilens, T. E., & Dodson, W. (2004). A clinical perspective of attention-deficit/hyperactivity disorder into adulthood. *Journal of Clinical Psychiatry*, 65, 1301-1313.
- Willcutt, E. G. (2015). Theories of ADHD. In R. A. Barkley (Ed.), *Attention-deficit hyperactivity disorder: A handbook for diagnosis and treatment* (4th ed.). New York, NY: Guilford.
- Willcutt, E. G., Doyle, A. E., Nigg, J. T., Faraone, S. V., & Pennington, B. F. (2005). Validity of the executive function theory for attention-deficit/hyperactivity disorder: A meta-analytic review. *Biological Psychiatry*, 57, 1336-1346.
- Willcutt, E. G., Betjemann, R. S., McGrath, L. M., Chhabildas, N. A., Olson, R. K., DeFries, J. C., & Pennington, B. F. (2010). Etiology and neuropsychology of comorbidity between RD and ADHD: The case for multiple-deficit models. *Cortex*, 46, 1345-1361.
- Willcutt, E. G., Nigg, J. T., Pennington, B. F., Solanto, M. V., Rohde, L. A., Tannock, R., ... Lahey, B. B. (2012). Validity of DSM-IV attention deficit/hyperactivity disorder symptom dimensions and subtypes. *Journal of Abnormal Psychology*, 121, 991-1010.
- Willcutt, E. G., Pennington, B. F., Boada, R., Ogline, J. S., Tunick, R. A., Chhabildas, N. A., & Olson, R. K. (2001). A comparison of the cognitive deficits in reading disability and attention-deficit/hyperactivity disorder. *Journal of Abnormal Psychology*, 110, 157-172.
- Williams, D. M., Dunsiger, S., Ciccolo, J. T., Lewis, B. A., Albrecht, A. E., & Marcus, B. H. (2008). Acute affective response to a moderate-intensity exercise stimulus predicts physical activity participation 6 and 12 months later. *Psychology of Sport and Exercise*, 9, 231-245.
- Winter, B., Breitenstein, C., Mooren, F. C., Voelker, K., Fobker, M., Lechtermann, A., ... Knecht, S. (2007). High impact running improves learning. *Neurobiology of Learning and Memory*, 87, 597-609.
- Woods, S. P., Lovejoy, D. W., & Ball, J. D. (2002). Neuropsychological characteristics of adults with ADHD: A comprehensive review of initial studies. *Clinical Neuropsychologist*, 16, 12-34.
- Zylowska, L., Ackerman, D. L., Yang, M. H., Futrell, J. L., Horton, N. L., Hale, T. S., et al. (2008). Mindfulness meditation training in adults and adolescents with ADHD: A feasibility study. *Journal of Attention Disorders*, 11, 737-746.